

## **Health & Mental Health**

- The Correctional Investigator reports that health care is the area of concern most frequently identified by prisoners in Canada.<sup>1</sup>
- HIV/AIDS was recognized as an international security issue by the United Nations Security Council in January 2000.
- Throughout Canada and the world, rates of HIV and Hepatitis C infection in prison populations are much higher than those found in the general population.<sup>2</sup>
- The HIV infection rate among women in prison is significantly higher than among male prisoners (3.6% v. 1.6%).<sup>3</sup>
- In a 2003 study, reported rates of hepatitis C infection were also much higher among women prisoners than among men prisoners (29.2% v. 16.6%).<sup>4</sup>
- Prisoners are 30 times more likely to have been infected with Hepatitis C and 7 to 10 times more likely to be infected with HIV than the general Canadian population.<sup>5</sup>
- Since most prisoners will eventually be released into the community, infection rates pose a threat to public health as well as the health of those who are currently imprisoned.<sup>6</sup>
- Women represent an increasing proportion of reported HIV cases in Canada. The greatest increase in new infections has been among young women, aged 15 to 29 years. At present, heterosexual transmission accounts for nearly 75% of all new infections in women.<sup>7</sup>
- While women as a group are more vulnerable than men to HIV infection and AIDS-related illnesses, some populations of women face significantly greater risks. For example, HIV affects more than twice as many Aboriginal as non-Aboriginal women in Canada. As elsewhere in the world, women in Canada who are most disadvantaged and marginalized are also most vulnerable to HIV.<sup>8</sup>

---

<sup>1</sup> Howard Sapers, Annual Report of the Correctional Investigator, 2006-2007, (Ottawa: Minister of Public Works and Government Services Canada, 2007) at 37.

<sup>2</sup> Canadian HIV/AIDS Legal Network, "HIV and hepatitis C transmission in prison" (2008) Factsheet

<sup>3</sup> Canadian HIV/AIDS Legal Network, "Women in Prison" (2008) Fact sheet.

<sup>4</sup> *Ibid.*

<sup>5</sup> Howard Sapers, Annual Report of the Correctional Investigator, 2006-2007, (Ottawa: Minister of Public Works and Government Services Canada, 2007) at 12.

<sup>6</sup> Canadian HIV/AIDS Legal Network, "HIV and hepatitis C in prisons: A comprehensive strategy" (2008) Fact sheet.

<sup>7</sup> Barbara Clow, "HIV/AIDS on the rise for Canadian Women" (The Canadian Women's Health Network, 2005) online: <<http://www.cwhn.ca/resources/sti/opedHIV.html>>

<sup>8</sup> *Ibid.*

- Non-sterile injection drug practices and lack of access to safe injection approaches significantly increase risk for HIV infection.
- Despite the recognition of the potential benefits of harm reduction initiatives, a prisoner-based needle exchange program has yet to be introduced to curtail the spread of infectious diseases such as Hepatitis C and HIV.<sup>9</sup>
- Although providing health care for over 100 years, of the CSC health care sites that provided health care to prisoners in 2006, 52 per cent of the sites failed to be accredited, 38 per cent were accredited with conditions, and only 10 per cent were fully accredited. Two key factors that prevented accreditation included the inadequacy of the existing clinical governance structure and the absence of continuing professional education and training for health care staff. Accreditation for the remaining sites was placed on hold.<sup>10 11</sup>

## **Mental Health**

- Women's life experiences, such as family stress, menstruation, child birth, menopause, chronic diseases, et cetera, are too often treated as pathological and result in women being chemically restrained rather than supported in moving through key life phases.<sup>12</sup>
- Women access the mental health system more frequently, receive treatment more often, and have higher rates of hospitalization for psychiatric problems than do men.
- Mental health issues faced by federally sentenced women are considerable and tend to be different than those of their male counterparts. Many women are survivors of childhood and adult abuse, and the post-traumatic effects of that abuse often impacts their mental health. Many federally sentenced women are diagnosed with mental illness and women in federal penitentiaries have a higher rate of self-mutilation and attempted suicide than their male counterparts. In many cases women harm themselves primarily as a means of coping with the distress caused by incarceration.<sup>13</sup>
- Despite the reality that most women classified as maximum security prisoners, who have a mental or intellectual disability, are described by correctional authorities as not being capable of "managing" in general population, there is no significant statistical difference in the institutional adjustment of women with disabling mental health issues as compared to women with no such diagnosis.

---

<sup>9</sup> Howard Sapers, Annual Report of the Correctional Investigator, 2005-2006, (Ottawa: Minister of Public Works and Government Services Canada, 2006)

<sup>10</sup> *Ibid* at 8.

<sup>11</sup> The numbers have not changed according to the 2007 Report of the Correctional Investigator.

<sup>12</sup> *Ibid* at 3.

<sup>13</sup> See research by Jan Heney.

- The Correctional Service of Canada has failed to demonstrate that it meets its statutory obligation to provide essential mental health care and reasonable access to non-essential mental health care in accordance with "professionally accepted standards." Over the last decade, the number of mentally ill prisoners has doubled, yet the level of non-punitive mental health services for prisoners, especially those in specialized psychiatric hospitals has remained the same or diminished.<sup>14</sup>
- For women with mental health issues, poverty is often associated with increased risk of violence and abuse, yet mental health systems still emphasize bio-medical over social factors in women's lives.
- The legacies of colonization, such as residential schooling, have resulted in cultural discontinuity and oppression in Aboriginal communities that have been tied to high rates of depression, alcoholism, suicide, and violence against Aboriginal women. Between 1989 and 1993, Aboriginal women in Canada were more than three times more likely to commit suicide than were non-Aboriginal women.
- There is a need for further research to examine the under theorized link between mental illness and addictions to help provide effective programs and services for women.

### **Fetal Alcohol Spectrum Disorders**

- Fetal Alcohol Spectrum Disorder (FASD) was introduced to umbrella the associated diagnoses of Fetal Alcohol Syndrome and Alcohol-Related Effects, such as Alcohol-Related Birth Defects (ARBD) and Alcohol Related Neuro-Developmental Disorder (ARND). Increasingly, studies focusing on the social factors of FASD have critiqued the cultural, racial, gendered, socio-economic and classed nature of theories regarding FAS, FASE and FASD.
- Fetal Alcohol Syndrome was first identified in 1973 in the American medical literature. Diagnosis relies upon detection of facial abnormalities, growth deficiencies and central nervous system impairment presumed due to alcohol consumption during the first trimester of pregnancy. The range and severity of the impact of maternal alcohol use is related to variations in the timing of alcohol use, variations in the amount of alcohol use, malnutrition, poor overall health of the mother and many other contextual matters.<sup>15</sup>
- The moral construction of the risks related to drinking during pregnancy individualizes the social responsibility on women, and has even resulted in the criminalization of pregnant women and mothers in some areas of the US.

---

<sup>14</sup> Howard Sapers, Annual Report of the Correctional Investigator, 2006-2007, (Ottawa: Minister of Public Works and Government Services Canada, 2007) at 23.

<sup>15</sup> Nancy Poole, "Mother and Child Reunion: Preventing Fetal Alcohol Spectrum Disorder by Promoting Women's Health" (BC Centre of Excellence for Women's Health, 2003) at 1, 2.

- In Canada, the dominant approach to preventing Fetal Alcohol Spectrum Disorders has been to focus on a single determinant: alcohol use. There must be further consideration of other situational and social risk factors that are related to FAS/E such as socioeconomic status, multiple non-prescription and prescription drug use, poor water and food quality and inadequate health care.<sup>16</sup>
- Prevention of FAS is largely based on this assumption of individualized responsibility and focuses only on substance abstinence rather than considering a need for collective action, attempts to ameliorate social inequality, attempt to ameliorate social inequality, inadequate health care, inadequate maternal nutrition, nor the impact of environmental pollutants et cetera on a fetus or a young child.<sup>17</sup>

### Addictions

- In a Saskatchewan study, treatment centre staff ranked lost cultural identity as the single most important factor for drug and alcohol abuse among Aboriginal people.
- Substance use among women in prison is estimated at over 90%. With the evisceration of publicly funded addiction, counseling and mental health services in the community. Prisons are increasingly serving as the default option to the lack of treatment centres.<sup>18</sup>
- Women's addictions differ from men by the type of drugs used, their social background and the reason for use, which underscores the need for gender specific substance use programs.
- Most criminalized women with substance use issues have significant histories of personal trauma and abuse, be it physical or sexual. Women frequently report substance use as an anaesthetizing means of masking emotion in order to cope with ongoing or unresolved trauma and abuse.
- Benzodiazepines can impede cognitive functioning and side effects include depression, memory deficiency, heightened emotions, and suicidal tendencies.<sup>19</sup> In Canada, the prescription of benzodiazepines increased by 12.8% from 1996 to 2000. Sleeping pill prescriptions have noticeably increased by 57.5%. Canadian and international studies have reported that 20 to 50% of women over the age of 60 have long term prescriptions to benzodiazepines and sleeping pills.<sup>20</sup> In western Canada, one out of

---

<sup>16</sup> *Ibid.*

<sup>17</sup> *Ibid* at 1.

<sup>18</sup> Shoshana Pollock, *Locked In, Locked Out: Imprisoning Women in the Shrinking and Punitive Welfare State*. (2008) at 14.

<sup>19</sup> Janet Currie, "Manufacturing Addiction: The Over-Prescription of Benzodiazepines and Sleeping Pills to Women in Canada", (BC Centre of Excellence for Women's Health, 2003) at 1.

<sup>20</sup> *Ibid* at 3.

three Aboriginal women over the age of 40 has been prescribed benzodiazepines.<sup>21</sup>

## **References**

Barbara Clow, “HIV/AIDS on the rise for Canadian Women” (The Canadian Women’s Health Network, 2005) online: <<http://www.cwhn.ca/resources/sti/opedHIV.html>>

Canadian HIV/AIDS Legal Network, “HIV and hepatitis C in prisons: A comprehensive strategy” (2008) Factsheet.

Canadian HIV/AIDS Legal Network, “HIV and hepatitis C transmission in prison” (2008) Factsheet.

Canadian HIV/AIDS Legal Network, “Women in Prison” (2008) Factsheet.

Canadian Human Rights Commission, *Protecting Their Rights: A Systemic Review of Human Rights in Correctional Services for Federally Sentenced Women* (2003).

Canadian Association of Social Workers, “The Declining Health and Well-Being of Low Income Women in Canada” (2006).

Canadian Institute for the Advancement of Women. *CRIAW Fact Sheet*. No. 6\_2004.

Correctional Service of Canada, *Infectious Diseases Prevention and Control in Canadian Federal Penitentiaries 2000-2001* (Ottawa: Correctional Service of Canada, 2003).

*Focus on Infectious Diseases: Health Care Needs of Inmates*, vol.3, no.1 (Ottawa: Correctional Service of Canada, 2005).

Health Canada, *Women and HIV/AIDS in Canada*, online: About Health Canada <[http://www.hc-sc.gc.ca/ahc-asc/pubs/int-aids-sida/womenfemmes\\_e.html](http://www.hc-sc.gc.ca/ahc-asc/pubs/int-aids-sida/womenfemmes_e.html)>

Howard Sapers, *Annual Report of the Office of the Correctional Investigator of Canada 2005-2006* (Ottawa: Minister of Public Works and Government Services Canada, 2006).

Howard Sapers, *Annual Report of the Correctional Investigator of Canada 2003-2004* (Ottawa: Minister of Public Works and Government Services Canada, 2004).

Janet Currie, “Manufacturing Addiction: The Over-Prescription of Benzodiazepines and Sleeping Pills to Women in Canada”, (BC Centre of Excellence for Women’s Health, 2003).

---

<sup>21</sup> *Ibid* at 4.

Lucy Hume, *A gender-specific substance abuse program for federally sentenced women* (Ottawa: Correctional Service of Canada, 2005).

Marina Morrow, "Mainstreaming Women's Mental Health: Building a Canadian Strategy" (2003) British Columbia Centre of Excellence for Women's Health

Nancy Poole, "*Mother and Child Reunion: Preventing Fetal Alcohol Spectrum Disorder by Promoting Women's Health*" (2003) British Columbia Centre of Excellence for Women's Health

Public Health Agency of Canada, *The Federal Initiative to Address HIV/AIDS in Canada: Strengthening Federal Action in the Canadian Response to HIV/AIDS* (Ottawa: Minister of Public Works and Government Services Canada, 2003).

Rick Lines *et al.*, *Prison Needle Exchange: Lessons from a Comprehensive Review of International Evidence and Experience*, online: Canadian HIV/AIDS Legal Network <<http://www.aidslaw.ca>>.

Shoshana Pollock, *Locked In, Locked Out: Imprisoning Women in the Shrinking and Punitive Welfare State*. (2008).